

Fibrous vein microstructure: Experimental and numerical simulation

C. Hilgers, J.L. Urai, A.D. Post & P.D. Bons

Geologie-Endogene Dynamik, RWTH Aachen, D-52056 Aachen

Department of Geological Science, Brown University, Providence RI 02912, USA

Department of Earth Sciences, Monash University, Clayton VIC 3168

Abstract

Syntectonic veins containing calcite or quartz grains with a pronounced fibrous morphology are thought to be formed by multiple crack-seal events, with the fibres growing parallel to the opening trajectory of the crack. This assumption provides a unique tool for structural geologists to analyse progressive deformation in rocks, although Cox (1987) and Williams and Urai (1989) have shown, that the long axis of the crystals do not always grow parallel to the opening direction. Urai et al. (1991) presented a model to explain the formation of fibrous morphologies, and predicted that, depending on the boundary conditions, fibres may or may not track the opening trajectory of the crack. We have further tested this model by simulations of natural antitaxial fibrous veins. Results show good agreement with observed morphology.

Introduction

One of the aims in structural geology is the reconstruction of the deformation history for small volumes of rock from the geometry and orientation of fabric elements (Lister & Williams, 1983). Such data can then be used for regional kinematic analysis. Methods for determination of finite strain and volume change are well established but deformation history is not so easily reconstructed. On the one hand progressive deformation overprints older structures, on the other hand all data must be retrieved from the final rock fabric. One of the few tools used in such analyses makes use of syntectonic fibrous veins (Beach, 1977; Ramsay & Huber 1983; Passchier & Trouw, 1995). Syntectonic veins and pressure fringes containing mineral (calcite or quartz) grains with a pronounced fibrous morphology are common in rocks at low metamorphic grades and high fluid pressure (at higher temperatures the fibrous morphology becomes thermodynamically unstable and will be destroyed by recrystallization). Fibrous veins are exposed often together with veins of more equant, blocky infill morphologies. Such crystal morphologies are not expected for crystals growing from hydrothermal solution, and explaining their genesis has been attempted by several workers since the pioneering work of Taber (1916, 1918) and Mügge (1919, 1928, 1930). The key process leading to the fibrous morphologies is thought to be the crack-seal process (Ramsay 1980) and/or diffusional processes (Bons & Jessel 1997, Durney & Ramsay 1973, Taber 1916, 1918). A striking feature of many vein fibres is their markedly curved shape, without the crystal lattice having the corresponding curvature. This led Durney & Ramsay (1973) to propose that the curvature in these fibres was developed during crystal growth, with the fibre long axis of the fibrous crystal extending parallel to the opening vector. This means that fibrous veins have a unique capability to record progressive deformation histories. Therefore curved fibres have rapidly become an important tool for structural geologists (Ramsay & Huber 1983, Beutner & Diegel 1985; Ellis 1986; Passchier & Urai 1988, Fisher et al. 1994). Less attention has been paid to testing the tracking hypothesis itself, in spite of the classic Durney and Ramsay (1973) paper, where some exceptions were already noted in the cases of some fibres in pressure fringes and in 'stretched crystals'. This point was further corroborated by Cox & Etheridge (1983) and Cox (1987) who showed examples of veins where fibres clearly did not track the opening direction.

*Belgian Symposium on Structural Geology and Tectonics, Leuven, 18-21.9.1997
Aard. Meded., 1997*

related references

- BEUTNER, E.C. & DIEGEL, F.A. 1985. Determination of fold kinematics from syntectonic fibres in pressure shadows, Martinsburg slate, New Jersey. *American Journal of Science*, 285, 16-50.
- BONS, P.D. & JESSEL, M.W. 1997. Experimental simulation of the formation of fibrous veins by localised dissolution-precipitation. *Mineralogical Magazine*, 61, 53-63.
- BONS, P.D. 1997. Crack-seal vein textures - a numerical model. Paper presented at 1997 EUG meeting.
- COX, S. F. 1987. Antitaxial crack-seal vein microstructures and their relationship to displacement paths. *Journal of Structural Geology*, 9(7), 779-787.
- COX, S.F. & ETHERIDGE, M.A. 1983. Crack-seal fibre growth mechanisms and their significance in the development of oriented layer silicate microstructures. *Tectonophysics*, 92, 147-170.
- DURNEY, D.W. & RAMSAY, J.G. 1973. Incremental strains measured by syntectonic crystal growths. In: De Jong K.A. and Scholten R. (Eds.) *Gravity and tectonics*, Wiley NY, 67-96.
- ELLIS, M. 1986. The determination of progressive deformation histories from antitaxial syntectonic crystal fibres. *Journal of Structural Geology*, 8, 701-709.
- FISHER, D.M., BRANTLEY, S.L., LI, T. & MEANS, W.D. 1995. Experimental antitaxial growth of fibrous crystals. I: Technique, fiber character, and implications. Geological Society of America annual meeting 1995, abstracts with programs, A70-A71.
- LISTER, G.S. & WILLIAMS, P.F. 1983. The partitioning of flow in flowing rock masses. *Tectonophysics*, 92, 1-33.
- MEANS, W.D. & LI, T. 1995. Experimental antitaxial growth of fibrous crystals. II: Internal structures. Geological Society of America annual meeting 1995, abstracts with programs, A70.
- MÜGGE, O. 1919. Vorgänge in der Gesteinswelt und ihre Messung. *Nachrichten v. d. kgl. Ges. d. Wiss. zu Göttingen. Geschäftl. Mitt.*, 78-100.
- MÜGGE, O. 1928. Über die Entstehung faseriger Minerale und ihrer Aggregationsformen. *Neues Jahrb., Beib.* 58 A, 303-348.
- MÜGGE, O. 1930. Bewegungen von Porphyroblasten in Phylliten und ihre Messung. *Neues Jb. Mineral. Geol. Paläontol.*, 61, 469-510.
- PABST, A. 1931. 'Pressure shadows' and the measurement of the orientation in rocks. *Journal Mineralogical Society of America*, 16, 55-70.
- PASSCHIER, C.W. & TROUW, R.A.J. 1995. *Microtectonics*. Springer, Berlin, 289 pp.
- POST, A.D. 1989. Aangroei op substraat: uit vrije oplossing en in aders. unpubl. MSc thesis, University of Utrecht.
- RAMSAY, J.G. 1980. The crack-seal mechanism of rock deformation. *Nature*, 284, 135-139.
- RAMSAY, J.G. & HUBER, M.I. 1983. *The Techniques of Modern Structural Geology*, Vol. I. Academic Press, London, 307 pp.
- SCHWINNER, R. 1926. Kristallisation und gerichteter Druck. *Tschermaks Mineralogische und Petrologische Mitteilungen*, 37 (3-6), 219-225.
- TABER, S. 1916. The genesis of asbestos and asbestiform minerals. *Bull. Am. Inst. Min. Eng.*, 119, 1973-1998.
- TABER, S. 1918. The origin of veinlets in the Silurian and Devonian strata of New York. *Journal of Geology*, 26, 56-73.
- URAI, J.L., WILLIAMS, P.F., & van ROERMOND, H.L.M. 1991. Kinematics of crystal growth in syntectonic fibrous veins. *Journal of Structural Geology*, 13(7), 823-836.
- WILLIAMS, P.F. & URAI, J.L. 1989. Curved vein fibres: An alternative explanation. *Tectonophysics*, 158, 311-333.